



CO₂
Human
Emissions

Progress report on service elements for CO₂ Earth observation integration

Frédéric Chevallier

che-project.eu



Co-ordinated by
 ECMWF



CO₂ Human Emissions

D5.1 Progress report on service elements for CO₂ Earth observation integration

Dissemination Level:	Public
Author(s):	Frédéric Chevallier (CEA)
Date:	17/12/2019
Version:	1.0
Contractual Delivery Date:	dd/mm/yyyy
Work Package/ Task:	WP5/ T5.1
Document Owner:	Organisation
Contributors:	Organisations
Status:	Final



CO₂ Human Emissions

CHE: CO₂ Human Emissions Project

Coordination and Support Action (CSA)
H2020-EO-3-2017 Preparation for a European
capacity to monitor CO₂ anthropogenic emissions

Project Coordinator: Dr Gianpaolo Balsamo (ECMWF)
Project Start Date: 01/10/2017
Project Duration: 39 months

Published by the CHE Consortium

Contact:

ECMWF, Shinfield Park, Reading, RG2 9AX,
gianpaolo.balsamo@ecmwf.int



The CHE project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 776186.

AIRBUS



Table of Contents

1	Executive Summary	5
2	Introduction	5
2.1	Background.....	5
2.2	Scope of this deliverable	6
2.2.1	Objectives of this deliverable	6
2.2.2	Work performed in this deliverable	6
2.2.3	Deviations and counter measures	6
4	Recommendations for operational prototype	9
5	Research priorities	11
6	Conclusion	12
	Acknowledgements	12
7	References	12

Tables

Table 1:	Implementation priorities linked to the domain (global, regional, local) and stream for application in the prototype: Near Real Time (NRT) and re-analysis (RA). An estimate of the effort required is given in person months.....	9
Table 2:	Research priorities linked to the domain (global, regional, local) and stream for application in the prototype: Near Real Time (NRT) and re-analysis (RA). An estimate of the effort required is given in person months.....	11

1 Executive Summary

The present chapter puts relevant information from the Copernicus CO₂ Monitoring Mission Requirements Document and from the three reports of the Copernicus Expert group and of the Task forces in perspective in the context of the prototype system which is being designed in CHE. It successively discusses the satellite CO₂ retrievals, the satellite non-CO₂ observations, the ground-based remote sensing observations and the in situ and flask-sampling observations. It highlights the large research needs for the identification of the role of each relevant Earth observation type in the Copernicus CO₂ support capacity system and for the modelling capability associated to the main ones.

2 Introduction

2.1 Background

The CHE prototype aims at building a system to monitor the exchange of CO₂ and potentially other important man-made greenhouse gases like CH₄ between the Earth surface and the atmosphere with the use of observations (mostly in the atmosphere), models and prior information including the specification of their uncertainties. The system is designed to support the Paris Agreement and follows the directive of the EC as described by Task Forces on CO₂¹. The general rationale and strategy for the CHE prototype is provided in D5.9, stemming from the discussions in the first WP5 workshop (Reading, 25-26 September 2019). The main challenges will be explored with the following recommendations:

- **Multi-scale** approach to monitor emission from point sources (power stations or industrial facilities), cities and countries using different model domains from global, regional to local and model resolutions (e.g. from 25 km to 100 m).
- **Multi-species** approach to detect and attribute the observed atmospheric signal to specific sources/sinks (e.g. natural and anthropogenic emissions with sectoral distribution).
- **Multi-stream** approach to support different applications and users with a near-real time stream focusing on shorter synoptic timescales designed to provide early warnings and give feedback to data producers, and a re-analysis stream that uses consolidated quality-controlled data, products and models with their associated uncertainties to estimate trends.

Earth observation, the topic of this chapter, is the gathering of information about the physical, chemical and biological systems of the Earth by natural- and man-made- environment monitoring². The exploitation of Earth observations about atmospheric CO₂ will bring the primary added value of the Copernicus CO₂ support capacity compared to existing national greenhouse gas emission inventories that traditionally rely on activity data only. Helped in particular by unprecedented satellite imagery means, the CO₂ support capacity for anthropogenic CO₂ emissions will supply extra evidence on the emissions levels and trends (Pinty et al., 2017, p. 7) that will be merged or contrasted with existing knowledge. Its scope

¹ <https://www.copernicus.eu/en/news/news/new-co2-green-report-2019-published>

² https://www.earthobservations.org/g_faq.html

will not be limited to CO₂ satellite imagery and will cover many types of Earth observations that are related to CO₂ emissions or to CO₂ dispersion in the atmosphere. Together, the various Earth observation types will drive and support a complex emission-estimation process at various spatial scales from the very local one (a few hectares) to the global one: the list of potentially-useful data is exceptionally long. CHE is currently exploring a relatively small number of ways to complete CO₂ observations with other types of Earth observations: radiocarbon, NO₂, oxygen, solar-induced fluorescence, carbonyl sulfide, night-light intensity and fraction of absorbed photosynthetically active radiation in the plant canopy. However, at this early stage of development of an operational CO₂ support capacity with unprecedented ambition, it is important to keep many more strategies open, at least as second choices. In the end, they may all play some role in the operational system but some of them will be directly assimilated in the CO₂ system while some will only be used at the pre- or post-processing stage to better guide the Copernicus CO₂ support capacity or to characterize its skill. Weather observations form a typical example of this dilemma. Resolving it implies making choices on the modelling of uncertainty in the estimation problem (e.g., strong-constraint formulations vs. weak-constraint formulation of the data assimilation) that may dramatically affect the skill of the operational system.

The needs and requirements for Earth Observations in the future European CO₂ support capacity for anthropogenic CO₂ emissions have already been extensively discussed in a series of documents:

- The Copernicus CO₂ Monitoring Mission Requirements Document (Meijer et al., 2019)
- The three reports of the Copernicus Expert group and of the Task forces³

The present chapter does not aim at replacing or even paraphrasing those documents, but rather at putting their relevant information in perspective in the context of the prototype system which is being designed in CHE. It successively discusses the satellite CO₂ retrievals, the satellite non-CO₂ observations, the ground-based remote sensing observations and the in situ and flask-sampling observations.

2.2 Scope of this deliverable

2.2.1 Objectives of this deliverable

In this report we aim at reviewing and assessing the options of available observations for the CHE prototype.

2.2.2 Work performed in this deliverable

Synthesis of work performed in CHE WP1, WP3 and WP4, ESA MRD and Task Force CO₂ reports.

2.2.3 Deviations and counter measures

Not applicable

3 Earth Observation System components

3.1 Satellite CO₂ observations

³ <https://www.copernicus.eu/en/news/news/new-co2-green-report-2019-published>

A Copernicus CO₂ monitoring (CO₂M) constellation with imaging capability as described in Pinty et al. (2017) is considered a prerequisite for the success of the CO₂ support capacity (Pinty et al. 2019, p. 3), while the need for a strong ground-based infrastructure comes in addition to the Copernicus constellation (Meijer et al., 2019, p. 11). This chapter is therefore built with the assumption that this Copernicus constellation will be deployed in orbit. Otherwise, the emphasis on the various observation types would be different.

An extensive CO₂ plume imaging capacity will be the best asset of the future Copernicus CO₂ monitoring constellation for the monitoring of CO₂ anthropogenic emissions compared to existing satellite missions. It will be extensive spatially because of the large swath of each space-borne instrument (better than 250 km, requirement S7MR-OBS-010 in Meijer et al., 2019) joined with its high spatial resolution (better than 4 km², requirement S7MR-OBS-020 in Meijer et al., 2019). It will be extensive temporally too because the observing system will include copies of the same instrument deployed on satellites with different orbital characteristics.

The CO₂ plume imaging capacity will rely on the near-contiguous sampling of backscattered solar light in selected spectral bands within the swath of the instrument and along the track of the satellite. When observation conditions are favourable (which mainly means enough insolation with low cloud and aerosol contamination), the column average dry-air mole fraction of CO₂ (and associated vertical averaging kernel) will be retrieved at each viewing location with low systematic and random errors (requirements S7MR-DAT-010 and S7MR-DAT-050 in Meijer et al., 2019). This will allow for resolving CO₂ plumes from emission hot spots and their surroundings. The wealth of high-quality column retrievals will also allow constraining large-scale carbon budgets over the globe to an unprecedented level.

The Sentinel CO₂ constellation will be operated within a larger constellation of CO₂ sounders of various types and operated by several agencies (CEOS Atmospheric Composition Virtual Constellation Greenhouse Gas Team, 2018), which can help filling gaps between Sentinel orbits and characterizing the actual Sentinel CO₂ retrieval noise over time.

3.2 Satellite non-CO₂ observations

Satisfying the ambitious objectives of the European CO₂ support capacity for anthropogenic CO₂ emissions implies exploiting complementary Earth observations, including some from satellites.

Meijer et al. (2019) plan for aerosol and cloud information to be provided by different types of radiometers on-board the same platform and at the same location as the Sentinel CO₂ column retrievals. Such data will both help disentangling the CO₂ signal from the cloud and aerosol signals in the measured spectra, and help excluding the pixels where the signal remains too ambiguous.

Further in the data flow, NO₂ retrievals that are spatially and temporally co-located with the Sentinel CO₂ retrievals will allow some tagging of the CO₂ plumes with respect to cleaner “background” scenes to identify the CO₂ source and to characterize the plume direction and the local wind speed (Meijer et al., 2019, p. 24). NO₂ was chosen because it is co-emitted with CO₂ when fossil fuel is burnt; but, on the downside, CO₂ and NO₂ plumes may not always overlap, even close to the plume origin, because NO₂ has a lifetime of the order of hours while CO₂ does not have any defined lifetime.

Information about wind direction from NO₂ plumes will be restricted to the vicinity of NO₂ emission hot spots. It will have to be complemented by information about wind speed at the same location and by information about the 3D structure of wind over the whole globe when

inferring large-scale CO₂ budgets. This information can come from wind-dedicated satellites in particular and from a much larger range of weather observations assimilated in Numerical Weather Prediction systems in general.

Many satellite observations can also provide some valuable information related to CO₂ sources and sinks in vegetated areas (solar-induced fluorescence, green fraction of absorbed photosynthetically active radiation in the plant canopy, vegetation biomass, ...) or about CO₂ anthropogenic emissions (NO₂⁴, CO, ...). Some other satellite observations can serve as proxies for the spatial and temporal variations of CO₂ emissions (data from the Global Positioning System, night light imagery, ...).

3.3 Ground-based remote sensing observations

The above-mentioned satellite retrievals of CO₂ and NO₂ columns and of aerosol properties are traditionally tuned⁵ with the help of ground-based radiometers that also observe vertically-integrated quantities. Reference retrievals from such devices are organized in international networks, like the Total Carbon Column Observing Network (TCCON, Wunch et al., 2011) and the Collaborative Carbon Column Observing Network (COCCON, Frey et al., 2019) for CO₂ or the Aerosol Robotic Network (AERONET, Holben et al., 1998) for aerosols. However, the requirement on systematic errors for the CO₂ column in the support capacity for anthropogenic CO₂ emissions (better than 0.5 ppm, requirement S7MR-DAT-050 in Meijer et al., 2019) is not less stringent than that of the current TCCON retrievals (Wunch et al., 2015) and will have to be improved for them to play a reference role here. Also, the spatial coverage of the CO₂ observations remains very sparse.

3.4 In situ and flask-sampling observation

Article 3 of the Copernicus regulation⁶ defines “in situ data” as all Earth observation data and ancillary ones that are not made from space. We choose the standard English definition here: “in the original or correct place”⁷, that excludes ground-based remote-sensing and flask samples analysed in distant laboratories. We therefore distinguish between in situ observations and flask-sampling observations in this section and we separate this section from the previous one about ground-based remote-sensing observations.

Relevant observations for the European CO₂ support capacity obviously include observations of the target quantities that can be used at the minimum for validation: local flux observations from micrometeorological towers in urban or other environments (like the Urban Flux Network or FluxNet), or from stack emission monitoring systems. They also cover local observations that either can be directly assimilated, or can inform about the quality of the assimilated data, or can help numerical models used in the data assimilation process. The more observation types and model types involved in the European CO₂ support capacity, the more Earth observations will be needed to support them. For example, satellite retrievals need observations for tuning; any additional tracer observation that is assimilated needs information about its sources and sinks and about how it relates to CO₂ emissions; any numerical model used in the assimilation process requires uncertainty quantification. It is important to note that there will be nuisance variables (i.e. knowledge gaps that are not limited to the target variables

⁴ NO₂ was mentioned above for plume tagging. Here, we mention its use for data assimilation.

⁵ Following the definition of BIPM (<https://www.bipm.org/fr/publications/guides/vim.html>), we do not use the word “calibrated” here that implies controlled conditions that are not possible in the open air.

⁶ <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX%3A32014R0377>

⁷ <https://www.oxfordlearnersdictionaries.com/definition/english/in-situ>

of the Copernicus support capacity) for all atmospheric tracers, even for the most obvious ones here like radiocarbon (Wang, 2016).

Given the emphasis put on atmospheric dispersion in the European CO₂ support capacity for anthropogenic CO₂ emissions, it is natural to quote in situ and flask-sampling tracer observations first. They are made close to the Earth's surface or higher in the atmosphere by aircraft, balloons or from free-fall tubes. They include observations of the main tracers mentioned so far in this chapter (CO₂, NO₂, CO), but also of isotopic measurements like radiocarbon, for other tracers that are related to CO₂ sources and sinks (e.g., oxygen and carbonyl sulfide) or to atmospheric transport (e.g., radon and SF₆). Reference measurements of that kind are maintained within programmes coordinated by the Global Atmospheric Watch Programme of the World Meteorological Organization (like those of the European Integrated Carbon Observation System, or of the National Oceanic and Atmospheric Administration Greenhouse Gas Reference Network, or the long-term aircraft programmes Comprehensive Observation Network for TRace gases by AirLiner and In-service Aircraft for a Global Observing System). Urban networks of lower-cost medium precision sensors for greenhouse gases may also be deployed in the future (e.g., Wu et al., 2016). Some data are available from air quality networks. Some come with delays that are not fit for near-real-time data assimilation and leaving them suitable for post-processing (e.g., validation) or re-analyses. Aircore measurements of the CO₂ mole fraction profile (Tans, 2009) have a special role in this domain because they are the only calibrated measurements of the CO₂ column per se and therefore come with very small systematic errors. However, they cannot be operated in an urban (i.e. inhabited) environment for security reasons. We also mention here again the need for accurate information about atmospheric winds when inferring emissions from mole fraction gradients, and the interest of exploiting proxy observations for the spatial and temporal variations of CO₂ emissions, like road traffic or temperature data. In terms of emission model support, the main need will likely be for observations informing about emission factors or emission ratios which vary much in space and time (e.g., Ammoura et al., 2014).

In situ and flask-sampling observation all have heterogeneous spatial coverage, in particular outside developed countries. They therefore do not sample the natural variability of their target variables well.

4 Recommendations for operational prototype

As explained in the introduction, it is still too early to define the role of each Earth observation stream in the operational prototype. Only the data from the CO₂M mission have a clear position in the system, as assimilated data. The measurement systems or measurement networks for the other data that will be given a key role will likely need to be developed, but this is not addressed here.

The above-mentioned Earth observations can be summarised in the following form.

Table 1: Implementation priorities linked to the domain (global, regional, local) and stream for application in the prototype: Near Real Time (NRT) and re-analysis (RA). An estimate of the effort required is given in person months.

Component	Domain	Stream	Recommendation	Estimated effort (Person Months)
Satellite XCO ₂ retrievals (Atmospheric Composition)	Global, regional	NRT, RA	Timeliness, accuracy	60

CO₂ HUMAN EMISSIONS 2019

Virtual Constellation including the CO ₂ M mission)				
Satellite retrievals of column-average non-CO ₂ tracers related to CO ₂ anthropogenic emissions (NO ₂ , CO, ...),	Global, regional	NRT, RA	Timeliness, accuracy	60
Satellite retrievals related to CO ₂ sources and sinks in vegetated areas (SIF, FAPAR, vegetation biomass, ...)	Global, regional	NRT, RA	Timeliness, accuracy	60
Satellite observations related to the spatial and temporal variations of CO ₂ emissions (data from the GPS, night-light imagery, ...).	Global, regional	NRT, RA	Timeliness, accuracy	36
Ground-based remote sensing observations (TCCON, COCCON, AERONET)	Global, regional	RA	Accuracy (higher than current for XCO ₂)	6
In situ and flask-sampling observations of tracers that are related to CO ₂ sources and sinks or to atmospheric transport (WMO/GAW including ICOS, NOAA, IAGOS, CONTRAIL; air-quality networks and some lower-cost medium precision sensor urban networks)	Global, regional	NRT, RA	Timeliness, accuracy	36

Flux observations (FluxNet, the Urban Flux Network, ...)	Global, regional	RA	Accuracy	6
Wind observations from satellites or from the surface	Global, regional	NRT, RA	Timeliness, accuracy	60
Observations about sources and sinks of any non-CO ₂ tracer used and about how this tracer relates to CO ₂ emissions	Global, regional	NRT, RA	Timeliness, accuracy	60

5 Research priorities

The assimilation of Earth observations in an atmospheric chemistry-transport model, possibly coupled with emission and absorption process models, to monitor anthropogenic CO₂ emissions is a new promising research area. However, the ambition of the Copernicus CO₂ support capacity to reach enough accuracy in this domain for the provision of extra evidence on the anthropogenic emissions levels and trends within the framework of the Paris Agreement is particularly challenging. A large research effort is needed to identify the role of each relevant Earth observation type in such a system and to develop the modelling capability associated to the main ones. The priorities are summarised in the following table.

Table 2: Research priorities linked to the domain (global, regional, local) and stream for application in the prototype: Near Real Time (NRT) and re-analysis (RA). An estimate of the effort required is given in person months.

Component	Domain	Stream	Recommendation	Estimated effort (Person Months)
Transport model	Global	NRT, RA	Identify contributions from direct or indirect observations of atmospheric transport (plume orientation in the CO ₂ M mission images, wind vector retrievals, measurements of radon, ...) for the improvement of transport simulation	60

Data assimilation system	Global	NRT, RA	Identify contributions from direct or indirect observations of vegetation activity (SIF, COS, ...) for the separation between fossil fuel and non-fossil fuel fluxes	60
Data assimilation system	Global	NRT, RA	Identify contributions from direct or indirect observations of anthropogenic activity (co-emitted tracers, ...) for the separation between fossil fuel and non-fossil fuel fluxes and possibly for some sectoral attribution	60
Observation operator	Global	NRT, RA	Develop a realistic modelling framework for each new observation, including corresponding error statistics	60

6 Conclusion

In the final report, we will include the relevant conclusions from WP1-4 studies of the CHE project.

Acknowledgements

The author is grateful to Anna Agustí-Panareda, Marko Scholze, Thomas Kaminski, Jérôme Barré, Joe McNorton, Michael Vossbeck and Hartmut Boesch for constructive comments on a previous version of this report.

7 References

Ammoura, L., Xueref-Remy, I., Gros, V., Baudic, A., Bonsang, B., Petit, J.-E., Perrussel, O., Bonnaire, N., Sciare, J., and Chevallier, F.: Atmospheric measurements of ratios between CO₂ and co-emitted species from traffic: a tunnel study in the Paris megacity, *Atmos. Chem. Phys.*, 14, 12871–12882, <https://doi.org/10.5194/acp-14-12871-2014>, 2014.

CEOS Atmospheric Composition Virtual Constellation Greenhouse Gas Team: A constellation architecture for monitoring carbon dioxide and methane from space, Report from the Committee on Earth Observation Satellites (CEOS) Atmospheric Composition Virtual Constellation (AC-VC), available at:

http://ceos.org/document_management/Virtual_Constellations/ACC/Documents/CEOS_AC-VC_GHG_White_Paper_Version_1_20181009.pdf, 2018.

Frey, M., Sha, M. K., Hase, F., Kiel, M., Blumenstock, T., Harig, R., Surawicz, G., Deutscher, N. M., Shiomi, K., Franklin, J. E., Bösch, H., Chen, J., Grutter, M., Ohyama, H., Sun, Y., Butz, A., Mengistu Tsidu, G., Ene, D., Wunch, D., Cao, Z., Garcia, O., Ramonet, M., Vogel, F., and Orphal, J.: Building the COllaborative Carbon Column Observing Network (COCCON): long-term stability and ensemble performance of the EM27/SUN Fourier transform spectrometer, *Atmos. Meas. Tech.*, 12, 1513–1530, <https://doi.org/10.5194/amt-12-1513-2019>, 2019.

Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET: A federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16, 1998.

Meijer et al. (2019), Copernicus CO₂ Monitoring Mission Requirements Document, https://esamultimedia.esa.int/docs/EarthObservation/CO2M_MRD_v2.0_Issued20190927.pdf

Pinty, B., G. Janssens-Maenhout, M. Dowell, H. Zunker, T. Brunhes, P. Ciais, D. Dee, H. Denier van der Gon, H. Dolman, M. Drinkwater, R. Engelen, M. Heimann, K. Holmlund, R. Husband, A. Kentarchos, Y. Meijer, P. Palmer and M. Scholze (2017) An operational anthropogenic CO₂ emissions monitoring & verification support capacity - Baseline requirements, Model components and functional architecture, doi:10.2760/39384, European Commission Joint Research Centre, EUR 28736 EN.

Pinty B., P. Ciais, D. Dee, H. Dolman, M. Dowell, R. Engelen, K. Holmlund, G. Janssens-Maenhout, Y. Meijer, P. Palmer, M. Scholze, H. Denier van der Gon, M. Heimann, O. Juvyns, A. Kentarchos and H. Zunker (2019) An Operational Anthropogenic CO₂ Emissions Monitoring & Verification Support Capacity – Needs and high level requirements for in situ measurements, doi:10.2760/182790, European Commission Joint Research Centre, EUR 29817 EN.

Tans, P. P.: System and method for providing vertical profile measurements of atmospheric gases, US Patent Office, 2009.

Wang, Y. (2016) The potential of observations of radiocarbon in atmospheric CO₂ for the atmospheric inversion of fossil fuel CO₂ emission at regional scale. *Earth Sciences*. Université Paris-Saclay, 2016. English. (NNT : 2016SACLV113). (tel-01529200)

Wu, L., Broquet, G., Ciais, P., Bellassen, V., Vogel, F., Chevallier, F., Xueref-Remy, I., and Wang, Y.: What would dense atmospheric observation networks bring to the quantification of city CO₂ emissions?, *Atmos. Chem. Phys.*, 16, 7743–7771, <https://doi.org/10.5194/acp-16-7743-2016>, 2016.

Wunch, D., Wennberg, P. O., Toon, G. C., Connor, B. J., Fisher, B., Osterman, G. B., Frankenberg, C., Mandrake, L., O'Dell, C., Ahonen, P., Biraud, S. C., Castano, R., Cressie, N., Crisp, D., Deutscher, N. M., Eldering, A., Fisher, M. L., Griffith, D. W. T., Gunson, M., Heikkinen, P., Keppel-Aleks, G., Kyrö, E., Lindenmaier, R., Macatangay, R., Mendonca, J., Messerschmidt, J., Miller, C. E., Morino, I., Notholt, J., Oyafuso, F. A., Rettinger, M., Robinson, J., Roehl, C. M., Salawitch, R. J., Sherlock, V., Strong, K., Sussmann, R., Tanaka, T., Thompson, D. R., Uchino, O., Warneke, T., and Wofsy, S. C.: A method for

evaluating bias in global measurements of CO₂ total columns from space, *Atmos. Chem. Phys.*, 11, 12317–12337, <https://doi.org/10.5194/acp-11-12317-2011>, 2011.

Wunch, D., Toon, G. C., Sherlock, V., Deutscher, N. M., Liu, C., Feist, D. G., and Wennberg, P. O.: Documentation for the 2014 TCCON Data Release (Version GGG2014.R0), CaltechDATA, <https://doi.org/10.14291/tccon.ggg2014.documentation.r0/1221662>, 2015.

Document History

Version	Author(s)	Date	Changes
0.1	F. Chevallier (CEA/LSCE)	05/12/2019	Initial version
1.0	F. Chevallier (CEA/LSCE)	17/12/2019	Final version after internal review

Internal Review History

Internal Reviewers	Date	Comments
Michael Vossbeck and Thomas Kaminski (iLab)	17/12/2019	Approved with comments
Hartmut Boesch (UoL)	17/12/2019	Approved with comments

Estimated Effort Contribution per Partner

Partner	Effort
CEA	0.5
Total	0.5

This publication reflects the views only of the author, and the Commission cannot be held responsible for any use which may be made of the information contained therein.